

# Gefcin

Gemifloxacin 320 mg

## COMPOSITION

**GEFCIN 320 mg Tablet:** Each film coated tablet contains Gemifloxacin 320 mg as Gemifloxacin Mesylate INN.

## PHARMACOLOGY

Gemifloxacin is a fluoroquinolone antibiotic. It is bactericidal with minimum bactericidal concentrations. Gemifloxacin acts by inhibiting DNA synthesis through inhibition of the bacterial type II topoisomerases, DNA gyrase, and/or topoisomerase IV (TOPO IV) which are both essential for bacterial growth.

Gemifloxacin is rapidly absorbed after oral administration. It is widely distributed throughout the body. Studies in healthy subjects showed that Gemifloxacin is distributed rapidly into target tissues and body fluids such as the lung (epithelial lining fluid, alveolar macrophages, bronchial tissue) and nasal secretions.

Following oral administration of Gemifloxacin, approximately 36% and 61% of the dose is excreted in the urine and feces, respectively, as unchanged drug and metabolites. AUC values were generally only slightly higher (approx. 10%) in women than in men. No dose adjustment is required based on gender

## INDICATIONS

Gemifloxacin is indicated for the treatment of the following bacterial infections in adults caused by sensitive organisms as follows-

**Acute bacterial exacerbation of chronic bronchitis-** caused by *Streptococcus pneumoniae*, *Haemophilus influenzae*, *Haemophilus parainfluenzae*, or *Moraxella catarrhalis*.

**Community-acquired pneumonia (of mild to moderate severity)-** caused by *Streptococcus pneumoniae* (including multi-drug resistant strains [MDRSP])\* , *Haemophilus influenzae*, *Moraxella catarrhalis*, *Mycoplasma pneumoniae*, *Chlamydia pneumoniae*, or *Klebsiella pneumoniae*.

## DOSAGE AND ADMINISTRATION

GEFCIN can be taken with or without food and should be swallowed with a liberal amount of liquid. The recommended dose of GEFCIN<sup>TM</sup> is 320 mg daily, according to the following table:

INDICATION	DOSE	DURATION
Acute bacterial Exacerbation of Chronic Bronchitis (AECB)	One 320 mg tablet daily	5 Days
Community acquired pneumonia (CAP) mild to moderate severity		
due to known or suspected <i>S. pneumoniae</i> , <i>H. influenzae</i> , <i>M. pneumoniae</i> , or <i>C. pneumoniae</i> infection	One 320 mg tablet daily	5 Days
due to known or suspected MDRSP*, <i>K. pneumoniae</i> , or <i>M. catarrhalis</i> infection	One 320 mg tablet daily	7 Days

\*MDRSP: multi-drug resistant *Streptococcus pneumoniae*, includes isolates previously known as PRSP (penicillin-resistant *Streptococcus pneumoniae*), and are strains resistant to two or more of the following antibiotics: penicillin (MIC  $\geq 2$   $\mu\text{g/ml}$ ), 2nd generation cephalosporins (e.g., cefuroxime), macrolides, tetracyclines and trimethoprim/sulfamethoxazole.

## Patients with renal impairment

Dose adjustment in patients with mild/moderate renal impairment is not required. Some modification of dosage is recommended for patients with severe renal dysfunction. The following table provides dosage guidelines for use in patients with renal impairment:

Creatinine Clearance (mL/min)	Dose
$\geq 40$	See usual Dosage
$< 40$	160 mg once daily

Patients on haemodialysis or continuous ambulatory peritoneal dialysis therapy should receive 160 mg once daily.

**Hepatic impairment:** GEFCIN may be given to patients with hepatic impairment, with no requirement for dose adjustment.

**Elderly patients:** Dose adjustment is not required

### **SIDE EFFECT**

The general adverse events include abdominal pain, diarrhea, headache, nausea, rash and vomiting. Some side effects have been infrequently reported such as fungal overgrowth in body, dizziness and insomnia, urticaria, pruritis and a maculopapular erythematous skin rash.

### **SPECIAL PRECAUTION**

For patients with severe impairment of renal function, alteration of the dosage regimen to 160 mg once daily is necessary. Adequate hydration of patients receiving Gemifloxacin should be maintained to prevent the formation of a highly concentrated urine and crystalluria. Gemifloxacin may cause dizziness; if this occurs, patients should not operate an automobile or machinery or engage in activities requiring mental alertness or co-ordination.

### **USE IN PREGNANCY AND LACTATION**

Gemifloxacin should not be used in pregnant or lactating women. The safety and efficacy of Gemifloxacin in pregnant or lactating women have not been established.

### **CONTRAINDICATION**

Known hypersensitivity to Gemifloxacin and other quinolones, Patients who have previously suffered tendon damage with uroquinolones. Gemifloxacin should not be used in children under 18 years of age.

### **WARNING**

Tendinitis and tendon ruptures may occur in any age group during treatment with quinolones, including Gemifloxacin, but particularly in elderly patients or when corticosteroids are being co-administered. Gemifloxacin should be discontinued if tendinitis is suspected or at the first sign of pain or inflammation and the affected limb should be rested. In clinical studies with Gemifloxacin a small mean increase in QTc interval was observed. Gemifloxacin should be used with caution in patients predisposed to QTc interval prolongation or in patients taking other medications that are known to prolong the QTc interval. Gemifloxacin should be used with caution in patients with epilepsy.

### **INTERACTION**

Gemifloxacin absorption is significantly reduced when aluminium or magnesium containing antacids and iron salts are concomitantly administered. Gemifloxacin should be taken at least 2 hours before or 3 hours after these agents. Gemifloxacin should be taken at least 2 hours before sucralfate administration. No clinically significant interactions have been observed when Gemifloxacin was co-administered with omeprazole theophylline, digoxin, warfarin and oral contraceptives.

### **OVER DOSAGE**

No specific antidote is known. Dialysis does not remove Gemifloxacin sufficiently to be useful in overdose. In the event of acute oral overdosage, the stomach should be emptied by inducing vomiting or by gastric lavage; the patient should be carefully observed, treated symptomatically and adequate hydration should be maintained.

### **STORAGE CONDITION**

Store in a cool and dry place, protected from light and moisture. Keep out of reach of children.

### **COMMERCIAL PACK**

**GEFCIN 320 mg Tablet:** Box containing 2x4's tablet in Alu-Alu blister pack.

Manufactured by



For further query on the use of this medicine, consult to a registered Doctor or Pharmacist.